

RB Health (US) LLC

-PE Product Labels –

-continued

MAXIMUM STRENGTH

Mucinex[®] **SINUS**
NIGHTSHIFT

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

MAXIMUM STRENGTH NDC 72854-240-20

Mucinex[®]
NIGHTSHIFT

SINUS

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Tripolidine HCl – Antihistamine

NIGHTTIME
RELIEF FOR A BETTER
MORNING

✓ COUGH ✓ FEVER ✓ SORE THROAT
✓ RUNNY NOSE ✓ SNEEZING
✓ NASAL CONGESTION

20 CAPLETS FOR AGES 12+

FAST RELEASE

ACTUAL SIZE

Questions?
1-888-MUCINEX (1-888-682-4639)
You may also report side effects to
this phone number.

HEALTH • HYGIENE • HOME
www.mucinex.com
Patients: www.rb.com/patients

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health
3176508 011921

3 63824 26271 0

LOT:
EXP.:
MADE IN:
3176508



Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant
Triprolidine HCl 1.25 mg.....	Antihistamine

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
 - temporarily reduces fever
 - controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)**
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

Mucinex
NIGHTSHIFT

MAXIMUM STRENGTH

SINUS

Maximum Strength per 4-hour dose
Do not take more than a total of
8 caplets in any 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

MAXIMUM STRENGTH

Mucinex[®]
SINUS-MAX[®]

 **DAY**  **NIGHT**

VALUE PACK
24 LIQUID GELS

Tamper evident:
Do not use if carton is damaged or if printed
seal on blister is broken or missing.

Fast Dissolving Liquid Gels!

MAXIMUM STRENGTH

Mucinex[®]
SINUS-MAX[®]

 **DAY**  **NIGHT**

Acetaminophen – Pain Reliever
Dextromethorphan HBr - Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

✓ Relieves Sinus Pressure, Headache
& Congestion
✓ Controls Cough
✓ Thins & Loosens Mucus

Actual Size

Acetaminophen – Pain Reliever
Dextromethorphan HBr - Cough Suppressant
Doxylamine Succinate - Antihistamine
Phenylephrine HCl - Nasal Decongestant

✓ Relieves Nasal Congestion,
Sinus Pressure & Pain
✓ Controls Cough
✓ Relieves Runny Nose & Sneezing

Actual Size

16 LIQUID GELS
(Liquid Filled Capsules)

TOTAL 24 LIQUID GELS

8 LIQUID GELS
(Liquid Filled Capsules)

LOT:
EXP.:
3176793



Questions?
1-888-MUCINEX (1-888-622-4639)
You may also report side effects to
this phone number.


HEALTH • HYGIENE • HOME
www.mucinex.com
Patient: www.rh.com/patient

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health

3176793 031521



3 63824 68824 4

Drug Facts**Active ingredients (in each liquid gel) Purposes**
Mucinex SINUS-MAX DAY

Acetaminophen 325 mg.....Pain reliever
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Guaifenesin 200 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Active ingredients (in each liquid gel) Purposes
Mucinex SINUS-MAX NIGHT

Acetaminophen 325 mg.....Pain reliever
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Doxylamine succinate 6.25 mg.....Antihistamine
 Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - cough
 - minor aches and pains
 - sinus congestion and pressure
 - runny nose and sneezing (*NIGHT only*)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (*DAY only*)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (*NIGHT only*)

Drug Facts (continued)

■ a breathing problem such as emphysema or chronic bronchitis (*NIGHT only*)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*NIGHT only*)

When using this product ■ do not use more than directed

■ excitability may occur, especially in children (*NIGHT only*)

■ marked drowsiness may occur (*NIGHT only*)

■ alcohol, sedatives, and tranquilizers may increase drowsiness (*NIGHT only*)

■ avoid alcoholic drinks (*NIGHT only*)

■ be careful when driving a motor vehicle or operating machinery (*NIGHT only*)

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

■ cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see **Overdose warning**)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients (DAY only)

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Inactive ingredients (NIGHT only)

D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water



Mucinex
SINUS-MAX

MAXIMUM STRENGTH

Maximum Strength per 4-hour dose
 Do not take the Mucinex Sinus-Max DAY and NIGHT liquid gels at the same time. Always wait at least 4 hours before taking another dose of Mucinex Liquid gel. Do not take more than a total of 12 liquid gels in a 24-hour period.
 Take only as directed.
 Keep carton for full information

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org



Drug Facts

Active ingredients (in each caplet) Mucinex SINUS-MAX DAY

Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Purposes

Active ingredients (in each caplet) Mucinex SINUS-MAX NIGHT

Purposes

Acetaminophen 325 mg Pain reliever
Diphenhydramine HCl 12.5 mg Antihistamine/cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses
Mucinex SINUS-MAX DAY ■ temporarily relieves:
 ■ nasal congestion ■ headache
 ■ minor aches and pains ■ sinus congestion and pressure
 ■ temporarily promotes nasal and/or sinus drainage
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex SINUS-MAX NIGHT ■ temporarily relieves:
 ■ nasal congestion ■ headache
 ■ minor aches and pains ■ sinus congestion and pressure
 ■ runny nose ■ sneezing
 ■ cough
 ■ temporarily promotes nasal and/or sinus drainage

3054364 09211Z PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 12 caplets in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ with any other product containing diphenhydramine, even one used on the skin (**NIGHT only**)
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Ask a doctor before use if you have
 ■ liver disease ■ heart disease ■ diabetes
 ■ high blood pressure ■ thyroid disease
 ■ trouble urinating due to an enlarged prostate gland
 ■ glaucoma (**NIGHT only**)
 ■ a breathing problem such as emphysema or chronic bronchitis (**NIGHT only**)
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin
 ■ taking sedatives or tranquilizers (**NIGHT only**)

When using this product
 ■ do not use more than directed
 ■ excitability may occur, especially in children (**NIGHT only**)
 ■ marked drowsiness may occur (**NIGHT only**)
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT only**)
 ■ avoid alcoholic drinks (**NIGHT only**)
 ■ be careful when driving a motor vehicle or operating machinery (**NIGHT only**)

HINGE

Drug Facts (continued)

Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ do not take more than directed (see Overdose warning)
 ■ do not take more than 12 caplets in any 24-hour period
 ■ adults and children 12 years of age and over: take 2 caplets every 4 hours

Drug Facts (continued)

■ children under 12 years of age: do not use

Other information
 ■ store between 20-25°C (68-77°F)

Inactive ingredients (Mucinex SINUS-MAX DAY only)
 croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients (Mucinex SINUS-MAX NIGHT only)
 croscarmellose sodium, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, methacrylic acid – ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium bicarbonate, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)
 You may also report side effects to this phone number.

HINGE



Drug Facts

Active ingredients (in each caplet) Mucinex SINUS-MAX DAY

Acetaminophen 325 mg.....	Pain reliever
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Purposes

Active ingredients (in each caplet) Mucinex SINUS-MAX NIGHT

Purposes

Acetaminophen 325 mg.....Pain reliever
 Diphenhydramine HCl 12.5 mg.....Antihistamine/cough suppressant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

Mucinex SINUS-MAX DAY

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex SINUS-MAX NIGHT

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - cough

■ temporarily promotes nasal and/or sinus drainage

3054364_092117

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin (**NIGHT only**)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT only**)
- a breathing problem such as emphysema or chronic bronchitis (**NIGHT only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**NIGHT only**)

When using this product

- do not use more than directed
- excitability may occur, especially in children (**NIGHT only**)
- marked drowsiness may occur (**NIGHT only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT only**)
- avoid alcoholic drinks (**NIGHT only**)
- be careful when driving a motor vehicle or operating machinery (**NIGHT only**)

HINGE

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours

Drug Facts (continued)

- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients (Mucinex SINUS-MAX DAY only)

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients (Mucinex SINUS-MAX NIGHT only)

croscarmellose sodium, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, methacrylic acid – ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium bicarbonate, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

HINGE



Drug Facts

Active ingredients (in each caplet) Mucinex SINUS-MAX DAY

Durnee

Active ingredients (in each caplet)	Purposes
Acetaminophen 325 mg.....	Pain reliever
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Active ingredients (in each caplet)	Purposes
Acetaminophen 325 mg.....	Pain reliever
Diphenhydramine HCl 12.5 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

Mucinex SINUS-MAX DAY

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex SINUS-MAX NIGHT

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - cough
- temporarily promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount

8335612 101718

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

ADHESIVE AREA

Drug Facts (continued)

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin (**NIGHT only**)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT only**)
- a breathing problem such as emphysema or chronic bronchitis (**NIGHT only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**NIGHT only**)

When using this product

- do not use more than directed
- excitability may occur, especially in children (**NIGHT only**)
- marked drowsiness may occur (**NIGHT only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT only**)
- avoid alcoholic drinks (**NIGHT only**)
- be careful when driving a motor vehicle or operating machinery (**NIGHT only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or

ADHESIVE AREA

ADHESIVE AREA

Drug Facts (continued)

- lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for

Drug Facts (continued)

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients (Mucinex SINUS-MAX DAY only)

corn starch, croscopovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Inactive ingredients (Mucinex SINUS-MAX NIGHT only)

ADHESIVE AREA

ADHESIVE AREA

ADH

Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

corn starch, croscarmellose sodium, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, ferric oxide yellow, magnesium stearate, methacrylic acid-ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.


ADHES

ADHESIVE AREA



Drug Facts		Drug Facts (continued)	
Active ingredients (in each liquid gel)	Purposes	Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
Acetaminophen 325 mg	Pain reliever	When using this product do not use more than directed	
		Stop use and ask a doctor if	





Warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

PEEL HERE ▶

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F)
- dosing cup provided

Drug Facts (continued)

- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this





NDC 63824-287-22

VALUE
PACK

Children's Mucinex®

DAY TIME

Multi-Symptom Cold

Cough Suppressant
Expectorant
Nasal Decongestant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg
Phenylephrine HCl 2.5 mg

- ✓ **Stuffy Nose**
- ✓ **Cough**
- ✓ **Chest Congestion**
- ✓ **Breaks up Mucus**

Very Berry
Flavor Liquid

Age 4+

4 FL OZ (118 mL)

NIGHT TIME

Multi-Symptom Cold

Pain Reliever/Fever Reducer
Antihistamine/Cough Suppressant
Nasal Decongestant

Acetaminophen 325 mg
Diphenhydramine HCl 12.5 mg
Phenylephrine HCl 5 mg

- ✓ **Stuffy Nose**
- ✓ **Cough**
- ✓ **Runny Nose & Sneezing**
- ✓ **Fever & Sore Throat**

Mixed Berry
Flavor Liquid

Age 6+

4 FL OZ (118 mL)

8 FL OZ (236 mL) TOTAL

Day Time Multi-Symptom Cold – Age 4+

Drug Facts

Active ingredients (in each 5 mL)	Purposes
Dextromethorphan HBr 5 mg.....	Cough suppressant
Guaifenesin 100 mg.....	Expectorant
Phenylephrine HCl 2.5 mg.....	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

PRINT SAFE AREA

KO

Day Time Multi-Symptom Cold – Age 4+ (cont'd)

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 3 mg
- tamper evident: do not use if neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate

Night Time Multi-Symptom Cold – Age 6+

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg.....	Antihistamine/ cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Night Time Multi-Symptom Cold – Age 6+ (cont'd)

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

■ dosing cup provided

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

DO NOT GIVE THE FIRST DOSE OF NIGHT TIME MULTI-SYMPTOM COLD SOONER THAN 4 HOURS AFTER THE LAST DOSE OF DAY TIME MULTI-SYMPTOM COLD UNLESS DIRECTED BY A DOCTOR.

DO NOT GIVE BOTH PRODUCTS AT THE SAME TIME OR GIVE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

3-63824-90994-3

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - nasal congestion
 - sneezing
 - controls cough to help your child get to sleep
 - temporarily reduces fever
 - minor aches and pains
 - sore throat
 - runny nose
 - cough

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other drug containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness ▶

- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F)
- dosing cup provided
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

Keep cartons for full directions for use.



Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in England ©2014 RB 020614 8140737

PRINT SAFE AREA

KO

Drug Facts (continued)

Inactive ingredients
(MUCINEX NIGHTSHIFT SEVERE COLD & FLU)
ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), polyethylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Maximum Strength per 4-hour dose

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

*FAST-MAX COLD & FLU helps to relieve these symptoms day or night; NIGHTSHIFT SEVERE COLD & FLU helps to relieve these symptoms at night

Do not take Mucinex Fast-Max Cold & Flu and Mucinex Nightshift Severe Cold & Flu at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid. **TAKE ONLY AS DIRECTED.**

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

NDC 63824-505-26

DAY & NIGHT COLD & FLU PACK

MAXIMUM STRENGTH



COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectoant • Phenylephrine HCl – Nasal Decongestant



MAXIMUM STRENGTH



SEVERE COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant • Triprolidine HCl – Antihistamine



FOR AGES 12+

TWO – 6 FL OZ (180 mL) bottles TOTAL – 12 FL OZ (360 mL)

FOR AGES 12+

No coating

Made in England Dist. by: RB Health (US), Parsippany, NJ 07054-0224 ©2020 RB Health

Drug Facts (continued)

Active ingredients (in each 20 mL) Purposes

MUCINEX NIGHTSHIFT SEVERE COLD & FLU
Acetaminophen 660 mg Pain reliever/fever reducer
Dextromethorphan HBr 20 mg Cough suppressant
Phenylephrine HCl 10 mg Nasal decongestant
Triprolidine HCl 25 mg Antihistamine

Drug Facts

Active ingredients (in each 20 mL) Purposes

MUCINEX FAST-MAX COLD & FLU
Acetaminophen 660 mg Pain reliever/fever reducer
Dextromethorphan HBr 20 mg Cough suppressant
Guaifenesin 400 mg Expectoant
Phenylephrine HCl 10 mg Nasal decongestant

Drug Facts (continued)

- marked drowsiness may occur (Nightsift Severe Cold & Flu only)
- acid, sedatives, and tranquilizers may increase drowsiness (Nightsift Severe Cold & Flu only)
- avoid alcoholic drinks (Nightsift Severe Cold & Flu only)
- use caution when driving a motor vehicle or operating machinery (Nightsift Severe Cold & Flu only)

Stop use and ask doctor if:

- fever, sore throat, or sleepiness occur
- fever, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts.

These could be signs of a serious condition.

If frequent or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: taking more than the recommended dose (pretend) may cause liver damage in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

MUCINEX FAST-MAX COLD & FLU

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

Drug Facts (continued)

- marked drowsiness may occur (Nightsift Severe Cold & Flu only)
- acid, sedatives, and tranquilizers may increase drowsiness (Nightsift Severe Cold & Flu only)
- avoid alcoholic drinks (Nightsift Severe Cold & Flu only)
- use caution when driving a motor vehicle or operating machinery (Nightsift Severe Cold & Flu only)

Stop use and ask doctor if:

- fever, sore throat, or sleepiness occur
- fever, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts.

These could be signs of a serious condition.

If frequent or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: taking more than the recommended dose (pretend) may cause liver damage in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

MUCINEX FAST-MAX COLD & FLU

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product
- do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

do not use within 14 days after stopping product

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use

Do not use





Drug Facts (continued)

Inactive ingredients (Mucinex Sinus-Max Severe Congestion & Pain) anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum
*may contain this ingredient

Inactive ingredients (Mucinex Nightshift Sinus) ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Do not take Mucinex Sinus-Max Severe Congestion & Pain and Mucinex Nightshift Sinus at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid.
TAKE ONLY AS DIRECTED.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DAY & NIGHT

SINUS CONGESTION & PAIN RELIEF

MAXIMUM STRENGTH* NDC 63824-115-66

Mucinex

SINUS-MAX

SEVERE CONGESTION & PAIN

Acetaminophen – Pain Reliever
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ Clears Sinus Congestion
- ✓ Relieves Headache
- ✓ Thins & Loosens Mucus

Mucinex

NIGHTSHIFT

SINUS

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Triprolidine HCl – Antihistamine

NIGHTTIME RELIEF FOR A BETTER MORNING

✓ COUGH ✓ FEVER ✓ SORE THROAT
✓ RUNNY NOSE ✓ SNEEZING ✓ NASAL CONGESTION

FOR AGES 12+

TWO – 6 FL OZ (180 mL) bottles

TOTAL – 12 FL OZ (360 mL)

FOR AGES 12+

No coating

Dist. by: RB Health (US), Parsippany, NJ 07054-0224

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

Drug Facts Active ingredients (in each 20 mL) Purposes Mucinex Sinus-Max Severe Congestion & Pain Acetaminophen 650 mg.....Pain reliever/fever reducer Dextromethorphan HBr 20 mg.....Cough suppressant Phenylephrine HCl 10 mg.....Nasal decongestant Triprolidine HCl 2.5 mg.....Antihistamine	Drug Facts (continued) Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers (Nightshift Sinus only) When using this product ■ do not use more than directed ■ excitability may occur, especially in children (Nightshift Sinus only) ■ marked drowsiness may occur (Nightshift Sinus only) ■ alcohol, sedatives, and tranquilizers may increase drowsiness (Nightshift Sinus only) ■ avoid alcoholic drinks (Nightshift Sinus only) ■ use caution when driving a motor vehicle or operating machinery (Nightshift Sinus only) Stop use and ask a doctor if ■ nervousness, dizziness, or sleepiness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with fever, rash, or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Directions Mucinex Sinus-Max Severe Congestion & Pain ■ do not take more than directed (see Overdose warning) ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosing cup provided ■ do not use dosing cup with other products ■ dose as follows or as directed by a doctor ■ adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours ■ children under 12 years of age: do not use Mucinex Nightshift Sinus ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided ■ do not use dosing cup with other products ■ dose as follows or as directed by a doctor ■ adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours ■ children under 12 years of age: do not use	Other information ■ each 20 mL contains: sodium 12 mg (Sinus-Max Severe Congestion & Pain only) and sodium 16 mg (Nightshift Sinus only) ■ store at 20-25°C (68-77°F) ■ do not refrigerate
Drug Facts Active ingredients (in each 20 mL) Purposes Mucinex Sinus-Max Severe Congestion & Pain Acetaminophen 650 mg.....Pain reliever Guaifenesin 400 mg.....Expectorant Phenylephrine HCl 10 mg.....Nasal decongestant	Drug Facts (continued) Uses Mucinex Sinus-Max Severe Congestion & Pain ■ temporarily relieves: ■ nasal congestion ■ minor aches and pains ■ sinus congestion and pressure ■ temporarily promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive Mucinex Nightshift Sinus ■ temporarily relieves these common cold and flu symptoms: ■ nasal congestion ■ minor aches and pains ■ sore throat ■ headache ■ sinus congestion and pressure ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes due to hay fever ■ temporarily reduces fever ■ controls cough to help you get to sleep	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning (Nightshift Sinus only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, tea, dizziness, rash, nausea, or vomiting, consult a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ glaucoma (Nightshift Sinus only) ■ trouble urinating due to an enlarged prostate gland (Nightshift Sinus only) ■ a troubling problem such as emphysema or chronic bronchitis (Nightshift Sinus only) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus)	



A KICKSTART FOR YOUR DAY



Mucinex
FAST-MAX
KICKSTART

MAXIMUM STRENGTH



Mucinex
FAST-MAX
NIGHT-TIME

POWERFUL RELIEF FOR YOUR NIGHT

POWERFUL MULTI-SYMPTOM COLD & FLU RELIEF

ALL IN ONE SEVERE COLD & FLU RELIEF » DAY OR NIGHT

KICKSTART SEVERE COLD & FLU (cont'd)

Drug Facts (continued)

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)

***FAST-MAX KICKSTART SEVERE COLD & FLU helps to relieve these symptoms day or night; FAST-MAX NIGHT TIME SEVERE COLD & FLU helps to relieve these symptoms at night**

Do not take Mucinex Fast-Max Kickstart Severe Cold & Flu and Mucinex Fast-Max Night Time Severe Cold & Flu at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid.

TAKE ONLY AS DIRECTED

Maximum Strength per 4-hour dose



3 63824 99697 4




NIGHT-TIME SEVERE COLD & FLU (cont'd)

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (scl), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)

MENTHOL FLAVORED LIQUID

FOR AGES 12+

TWO - 6 FL OZ (180 mL) bottles

RELIEF FOR YOUR NIGHT

TOTAL - 12 FL OZ (360 mL)

FOR AGES 12+

POWERFUL RELIEF FOR YOUR NIGHT

+ A KICKSTART FOR YOUR DAY



reckitt

Dist. by: RB Health (US)
Parsippany, NJ 07054-8224
© 2023 RB Health
3259828 04/05/23

NIGHT TIME SEVERE COLD & FLU

Drug Facts

Active Ingredients (In each 20 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- liver disease
- heart disease
- thyroid disease
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

KICKSTART SEVERE COLD & FLU

Drug Facts

Active Ingredients (In each 20 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- liver disease
- heart disease
- thyroid disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

- Stop use and ask a doctor if
 - nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion, or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur

cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

MAXIMUM STRENGTH

NDC 72854-138-66

Mucinex®

FAST MAX®

KICKSTART
SEVERE COLD & FLU

Acetaminophen » Pain Reliever/Fever Reducer

Dextromethorphan HBr » Cough Suppressant

Guaifenesin » Expectorant

Phenylephrine HCl » Nasal Decongestant

**COMEBACK
WITH A
KICK**



MENTHOL FLAVORED LIQUID

6 FL OZ (180 mL)

FOR AGES 12+

3259831

030823





Drug Facts**Active ingredients
(in each liquid gel)****Mucinex FAST-MAX DAY Cold & Flu**

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Guaifenesin 200 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes**Active ingredients
(in each liquid gel)****Mucinex FAST-MAX NIGHT Cold & Flu**

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Doxylamine succinate 6.25 mg.....Antihistamine
 Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes**Uses**

- temporarily relieves these common cold and flu symptoms:
 - sinus congestion and pressure
 - cough
 - headache
 - sore throat
 - sneezing (NIGHT only)
 - itching of the nose or throat (NIGHT only)
 - itchy, watery eyes due to hay fever (NIGHT only)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (DAY only)
- controls cough to help you get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric,

Drug Facts (continued)

or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (NIGHT only)
- a breathing problem such as emphysema or chronic bronchitis (NIGHT only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (NIGHT only)

When using this product

- do not use more than directed
- excitability may occur, especially in children (NIGHT only)
- marked drowsiness may occur (NIGHT only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (NIGHT only)
- avoid alcoholic drinks (NIGHT only)
- be careful when driving a motor vehicle or operating machinery (NIGHT only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Drug Facts (continued)**Inactive ingredients (DAY only)**

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Inactive ingredients (NIGHT only)

D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

*DAY COLD & FLU helps to relieve these symptoms day or night; NIGHT COLD & FLU helps to relieve these symptoms at night

Maximum Strength per 4-hour dose

Do not take the Mucinex Fast-Max DAY COLD & FLU and Mucinex Fast-Max NIGHT COLD & FLU liquid gels at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid gels. Do not take more than a total of 12 liquid gels in any 24-hour period.

Take only as directed.

Keep carton for full information.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

MAXIMUM STRENGTH

Mucinex
FAST-MAX

DAY COLD & FLU
NIGHT COLD & FLU

10 LIQUID GELS

Fast Dissolving Liquid Gels!

MAXIMUM STRENGTH NDC 72854-144-10

Mucinex
FAST-MAX

DAY COLD & FLU
NIGHT COLD & FLU

Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Guafenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Doxylamine succinate - Antihistamine
Phenylephrine HCl - Nasal Decongestant

ALL IN ONE*

ALL IN ONE*

FOR AGES 12+

ACTUAL SIZE

ACTUAL SIZE

6 LIQUID GELS
(Liquid Filled Capsules)

TOTAL 10 LIQUID GELS

4 LIQUID GELS
(Liquid Filled Capsules)

reckitt

Dist. by: RB Health US
Rexbury, NJ 07054-0226
©2023 RB Health
www.mucinex.com
Patents:
www.mucinex.com/patents

Questions? 1-866-MUCINEX (1-866-682-4639)

013123

3265234

6382499731

Drug Facts

Active ingredients (in each liquid gel)

Mucinex FAST-MAX DAY Cold & Flu

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Purposes

Active ingredients (in each liquid gel)

Mucinex FAST-MAX NIGHT Cold & Flu

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - sinus congestion and pressure
 - cough
 - headache
 - sore throat
 - sneezing (NIGHT only)
 - itching of the nose or throat (NIGHT only)
 - itchy, watery eyes due to hay fever (NIGHT only)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (DAY only)
- controls cough to help you get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric,

Drug Facts (continued)

or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (NIGHT only)
- a breathing problem such as emphysema or chronic bronchitis (NIGHT only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (NIGHT only)

When using this product

- do not use more than directed
- excitability may occur, especially in children (NIGHT only)
- marked drowsiness may occur (NIGHT only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (NIGHT only)
- avoid alcoholic drinks (NIGHT only)
- be careful when driving a motor vehicle or operating machinery (NIGHT only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Drug Facts (continued)

Inactive ingredients (NIGHT only)

D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, polyurethane, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Drug Facts (continued)

Inactive ingredients (DAY only)

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, polyurethane, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Drug Facts (continued)
Inactive ingredients (Mucinex FREEFROM Cold & Flu Daytime)
 ammonium glycyrrhizate, anhydrous citric acid, cochineal carmine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Inactive ingredients (Mucinex FREEFROM Cold & Flu Nighttime)
 ammonium glycyrrhizate, anhydrous citric acid, cochineal carmine, edetate disodium, flavors, glycerin (soy), maltodextrin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Made in Mexico
 Dist. by: RB Health (US)
 Parsippany, NJ 07054-0224

Do not take Mucinex FREEFROM COLD & FLU DAYTIME and Mucinex FREEFROM COLD & FLU NIGHTTIME at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid.

TAKE ONLY AS DIRECTED.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

NDC 63824-940-26



COLD & FLU DAYTIME

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER
 DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
 GUAFENESIN – EXPECTORANT
 PHENYLEPHRINE HCl – NASAL DECONGESTANT

COLD & FLU NIGHTTIME

ACETAMINOPHEN – PAIN RELIEVER/FEVER REDUCER
 DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
 PHENYLEPHRINE HCl – NASAL DECONGESTANT
 TRIPROLIDINE HCl – ANTIHISTAMINE

— **FREEFROM** —
 ARTIFICIAL FLAVORS, COLORS, & DYES • ALCOHOL & SUGAR

ELDERBERRY & CHERRY NATURAL FLAVOR

FOR AGES 12+ TWO - 6 FL OZ (180 mL) BOTTLES TOTAL - 12 FL OZ (360 mL) FOR AGES 12+

No coating

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

<p>Drug Facts (continued)</p> <p>Active ingredients (in each 20 mL) Purposes Mucinex FREEFROM Cold & Flu Nighttime</p> <p>Acetaminophen 680 mg Pain reliever/fever reducer Dextromethorphan HBr 20 mg Cough suppressant Phenylephrine HCl 10 mg Nasal decongestant Triprolidine HCl 2.5 mg Antihistamine</p>	<p>Drug Facts (continued)</p> <p>Active ingredients (in each 20 mL) Purposes Mucinex FREEFROM Cold & Flu Daytime</p> <p>Acetaminophen 680 mg Pain reliever/fever reducer Dextromethorphan HBr 20 mg Cough suppressant Guafenesin 400 mg Expectorant Phenylephrine HCl 10 mg Nasal decongestant</p>	<p>Drug Facts (continued)</p> <p>Uses MUCINEX FREEFROM COLD & FLU DAYTIME</p> <ul style="list-style-type: none"> temporarily relieves these common cold and flu symptoms: <ul style="list-style-type: none"> cough nasal congestion sore throat headache sinus congestion and pressure temporarily reduces fever helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive <p>MUCINEX FREEFROM COLD & FLU NIGHTTIME</p> <ul style="list-style-type: none"> temporarily relieves these common cold and flu symptoms: <ul style="list-style-type: none"> cough nasal congestion sore throat headache sinus congestion and pressure runny nose sneezing itching of the nose or throat itchy, watery eyes due to hay fever temporarily reduces fever controls cough to help you get to sleep <p>Warnings</p> <p>User warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4000 mg in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure glaucoma trouble urinating due to an enlarged prostate gland a breathing problem such as emphysema or chronic bronchitis persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) <p>Other information</p> <ul style="list-style-type: none"> each 20 mL contains: sodium 12 mg (Daytime only) and sodium 12 mg (Nighttime only) store at 20°-25°C (68°-77°F) do not refrigerate
---	---	--



rb
 HEALTH • HYGIENE • HOME
 Questions?

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

MAXIMUM STRENGTH

Mucinex

SINUS-MAX NIGHTSHIFT

 **DAY**

PRESSURE, PAIN & COUGH

 **NIGHT**

SINUS

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

MAXIMUM STRENGTH

NDC 72854-243-20

Mucinex

SINUS-MAX NIGHTSHIFT

 **DAY**

PRESSURE, PAIN & COUGH

Acetaminophen - Pain Reliever
Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

- ✓ Relieves Sinus Pressure & Congestion
- ✓ Relieves Headache
- ✓ Controls Cough
- ✓ Thins & Loosens Mucus

12 CAPLETS

 **NIGHT**

SINUS

Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Phenylephrine HCl - Nasal Decongestant
Triprolidine HCl - Antihistamine

NIGHTTIME RELIEF FOR A BETTER MORNING

✓ COUGH ✓ FEVER ✓ SORE THROAT
✓ RUNNY NOSE ✓ SNEEZING
✓ NASAL CONGESTION

8 CAPLETS

FAST RELEASE

TOTAL 20 CAPLETS

PARENTS:

Learn about their medicine dose

www.StopTheMedicineAbuse.org

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

 **HEALTH • HYGIENE • HOME**

www.mucinex.com

Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health

3176362 012721

 6382426273 4

LOT:

EXP.:

MADE IN:

3176362

Drug Facts	
Active ingredients (in each caplet)	Purposes
Mucinex Sinus-Max Day Pressure, Pain & Cough	
Acetaminophen 325 mg	Pain reliever

010521

PEEL CORNER TO REVEAL COMPLETE DRUG FACTS AND INFORMATION

Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Active ingredients (in each caplet)

Mucinex Nightshift Night Sinus

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant
Tripolidine HCl 1.25 mg.....	Antihistamine

Purposes

Uses

Mucinex Sinus-Max Day Pressure, Pain & Cough

- temporarily relieves:
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - temporarily promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- headache
- cough

Mucinex Nightshift Night Sinus

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - sinus congestion and pressure
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
 - temporarily reduces fever
 - controls cough to help you get to sleep
- minor aches and pains
- sneezing
- runny nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning (Nightshift Night Sinus only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Reseal Area

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (*Nightshift Night Sinus only*)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (*Nightshift Night Sinus only*)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Nightshift Night Sinus only*)

When using this product

- do not use more than directed
- excitability may occur, especially in children (*Nightshift Night Sinus only*)
- marked drowsiness may occur (*Nightshift Night Sinus only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Nightshift Night Sinus only*)
- avoid alcoholic drinks (*Nightshift Night Sinus only*)
- use caution when driving a motor vehicle or operating machinery (*Nightshift Night Sinus only*)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

Hinge

Hinge

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Mucinex Sinus-Max Day Pressure, Pain & Cough

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Mucinex Nightshift Night Sinus

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients

Mucinex Sinus-Max Day Pressure, Pain & Cough

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients

Mucinex Nightshift Night Sinus

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



MIN PT SIZE &
NET WT (g) 5 FL OZ 150g
Drug Facts Table
Active Ingredient Table

Pediatrician Recommended!

NDC 63824-945-34

TRIPLE PACK INCLUDES:

2X **DAYTIME**

1X **NIGHTTIME**

FREEFROM

**MULTI-SYMPTOM
COLD, FLU
& SORE THROAT**

ACETAMINOPHEN — PAIN RELIEVER/FEVER REDUCER
DEXTROMETHORPHAN HBr — COUGH SUPPRESSANT
GUAFENESIN — EXPECTORANT
PHENYLEPHRINE HCl — NASAL DECONGESTANT

**MULTI-SYMPTOM
COLD & FLU
NIGHTTIME**

ACETAMINOPHEN — PAIN RELIEVER/FEVER REDUCER
DEXTROMETHORPHAN HBr — COUGH SUPPRESSANT
PHENYLEPHRINE HCl — NASAL DECONGESTANT
TRIPROlidine HCl — ANTIHISTAMINE

FREEFROM
ARTIFICIAL FLAVORS, COLORS & DYES • ALCOHOL & SUGAR

**ELDERBERRY
& CHERRY NATURAL FLAVOR**

THREE - 4 FL. OZ (118 mL) bottles

TOTAL - 12 FL OZ (354 mL)

AGES 6+ YRS.

MULTI-SYMPTOM COLD & FLU NIGHTTIME — Ages 6+ YRS. (cont'd)

DAYTIME MULTI-SYMPTOM COLD, FLU & SORE THROAT — Ages 6+ YRS.

DAYTIME MULTI-SYMPTOM COLD, FLU & SORE THROAT — Ages 6+ YRS. (cont'd)

Drug Facts (continued)

When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 4 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, cochineal carmine, edetate disodium, FD&C Blue #1, FD&C Yellow #6, hydroxypropyl methylcellulose, polyethylene glycol, purified water, sodium benzoate, xanthan gum

FREEFROM

NO ARTIFICIAL
COLOR OR DYES

NO ARTIFICIAL
FLAVORS

ALCOHOL
& SUGAR FREE

TRIPLE





Do not take the Mucinex Fast-Max Day Severe Congestion & Cough and Mucinex Nightshift Night Cold & Flu caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. TAKE ONLY AS DIRECTED.

Drug Facts

Active ingredients (in each caplet)

Mucinex Fast-Max Day Severe Congestion & Cough

Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Purposes

Active ingredients (in each caplet)

Purposes

Mucinex Nightshift Night Cold & Flu

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Triprolidine HCl 1.25 mg.....	Antihistamine

Uses

Mucinex Fast-Max Day Severe Congestion & Cough

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the impulse to cough to help you get to sleep
 - the intensity of coughing
 - nasal congestion due to a cold

Mucinex Nightshift Night Cold & Flu

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning (Nightshift Night Cold & Flu only): This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert (Nightshift Night Cold & Flu only): Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning (Nightshift Night Cold & Flu only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Reseal Area

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. **(Nightshift Night Cold & Flu only)**
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease **(Nightshift Night Cold & Flu only)**
- heart disease **(Fast-Max Day Severe Congestion & Cough only)**
- high blood pressure **(Fast-Max Day Severe Congestion & Cough only)**
- thyroid disease **(Fast-Max Day Severe Congestion & Cough only)**
- diabetes **(Fast-Max Day Severe Congestion & Cough only)**
- glaucoma **(Nightshift Night Cold & Flu only)**
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis **(Nightshift Night Cold & Flu only)**
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin **(Nightshift Night Cold & Flu only)**
- taking sedatives or tranquilizers **(Nightshift Night Cold & Flu only)**

When using this product

- do not use more than directed
- excitability may occur, especially in children **(Nightshift Night Cold & Flu only)**
- marked drowsiness may occur **(Nightshift Night Cold & Flu only)**
- alcohol, sedatives, and tranquilizers may increase drowsiness **(Nightshift Night Cold & Flu only)**
- avoid alcoholic drinks **(Nightshift Night Cold & Flu only)**
- use caution when driving a motor vehicle or operating machinery **(Nightshift Night Cold & Flu only)**

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days **(Nightshift Night Cold & Flu only)**
- fever gets worse or lasts more than 3 days **(Nightshift Night Cold & Flu only)**

3068

Hinge

Hinge

Drug Facts (continued)

- redness or swelling is present (**Nightshift Night Cold & Flu only**)
- new symptoms occur (**Nightshift Night Cold & Flu only**)
- cough comes back, or occurs with fever, rash, or headache that lasts.
These could be signs of a serious condition. (**Nightshift Night Cold & Flu only**)
- nervousness, dizziness, or sleeplessness occur (**Fast-Max Day Severe Congestion & Cough only**)
- symptoms do not get better within 7 days or occur with fever
(**Fast-Max Day Severe Congestion & Cough only**)
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts.
These could be signs of a serious condition. (**Fast-Max Day Severe Congestion & Cough only**)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Overdose warning (Nightshift Night Cold & Flu only): Taking more than the recommended dose (overdose) may cause liver damage. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions**Mucinex Fast-Max Day Severe Congestion & Cough**

- do not take more than directed
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Mucinex Nightshift Night Cold & Flu

- do not take more than directed (see **Overdose warning**)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information ■ store at 20-25°C (68-77°F)**Inactive ingredients Mucinex Fast-Max Day Severe Congestion & Cough**

croscarmellose sodium, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, methacrylic acid-ethyl acrylate copolymer, mica, microcrystalline cellulose, polyethylene glycol 3350, polysorbate 80, polyvinyl alcohol, povidone K29/32, sodium bicarbonate, talc, titanium dioxide

Inactive ingredients Mucinex Nightshift Night Cold & Flu

croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.

Drug Facts (continued)

Other information

- each 20 mL contains: sodium 9 mg (Severe Congestion & Pain Clear & Cool only) and sodium 16 mg (Nightshift Sinus Clear & Cool only)
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients (Mucinex Sinus-Max Severe Congestion & Pain Clear & Cool) anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Inactive ingredients (Mucinex Nightshift Sinus Clear & Cool) ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, water, xanthan gum

Do not take Mucinex Sinus-Max Severe Congestion & Pain Clear & Cool and Mucinex Nightshift Sinus Clear & Cool at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid. TAKE ONLY AS DIRECTED.

MAXIMUM STRENGTH* NDC 63824-116-66

Mucinex[®]

SINUS-MAX[®]

SEVERE CONGESTION & PAIN

CLEAR & COOL[®]

Acetaminophen – Pain Reliever
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ **Clears Sinus Congestion**
- ✓ **Relieves Headache**
- ✓ **Thins & Loosens Mucus**

FOR AGES 12+ TWO – 6 FL OZ (180 mL) bottles

Mucinex[®]

NIGHTSHIFT

SINUS CLEAR & COOL

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Tripolidine HCl – Antihistamine

NIGHTTIME
RELIEF FOR A BETTER
MORNING

✓ COUGH ✓ FEVER ✓ SORE THROAT
 ✓ RUNNY NOSE ✓ SNEEZING ✓ NASAL CONGESTION

FOR AGES 12+ TOTAL – 12 FL OZ (360 mL)

Drug Facts	Drug Facts (continued)
Active ingredients (in each 20 mL) Purposes	Active ingredients (in each 20 mL) Purposes
Mucinex Sinus-Max	Mucinex Nightshift
SEVERE CONGESTION & PAIN CLEAR & COOL	Sinus Clear & Cool
Acetaminophen 650 mg.....Pain reliever	Acetaminophen 650 mg.....Pain reliever/fever reducer
Guaifenesin 400 mg.....Expectorant	Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant	Phenylephrine HCl 10 mg.....Nasal decongestant
	Tripolidine HCl 2.5 mg.....Antihistamine

Drug Facts (continued)

Uses **Mucinex Sinus-Max SEVERE CONGESTION & PAIN CLEAR & COOL**

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex Nightshift Sinus Clear & Cool

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning (Nightshift Sinus Clear & Cool only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (**Nightshift Sinus Clear & Cool only**)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (**Nightshift Sinus Clear & Cool only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nightshift Sinus Clear & Cool only**)

When using this product

- do not use more than directed
- excitability may occur, especially in children (**Nightshift Sinus Clear & Cool only**)
- marked drowsiness may occur (**Nightshift Sinus Clear & Cool only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nightshift Sinus Clear & Cool only**)
- avoid alcoholic drinks (**Nightshift Sinus Clear & Cool only**)
- use caution when driving a motor vehicle or operating machinery (**Nightshift Sinus Clear & Cool only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

MUCINEX SINUS-MAX SEVERE CONGESTION & PAIN CLEAR & COOL

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

MUCINEX NIGHTSHIFT SINUS CLEAR & COOL

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

MAXIMUM STRENGTH

Mucinex[®]

FAST-MAX[®]

COLD & FLU
ARCTIC BURST™

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant



FOR AGES 12+ TWO – 6 FL OZ (180 mL) bottles

NDC 72854-135-26

Mucinex[®]

NIGHTSHIFT™

SEVERE COLD & FLU
ARCTIC BURST™

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Triprolidine HCl – Antihistamine



NIGHTTIME
RELIEF FOR A BETTER
MORNING

✓ COUGH ✓ FEVER ✓ SORE THROAT
✓ RUNNY NOSE ✓ SNEEZING ✓ NASAL CONGESTION

TOTAL – 12 FL OZ (360 mL) FOR AGES 12+

Drug Facts (continued)**Other information**

- each 20 mL contains:
sodium 9 mg (Fast-Max
Cold & Flu Arctic Burst
only) and sodium 16 mg
(Nightshift Severe Cold &
Flu Arctic Burst only)
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients**(MUCINEX FAST-MAX COLD
& FLU ARCTIC BURST)**

anhydrous citric acid, D&C
yellow no. 10, edetate
disodium, FD&C blue no. 1,
flavors, glycerin, propyl gallate,
propylene glycol, sodium
benzoate, sodium citrate,
sorbitol, sucralose, water,
xanthan gum

Inactive ingredients**(MUCINEX NIGHTSHIFT SEVERE
COLD & FLU ARCTIC BURST)**

ammonium glycyrrhizate,
anhydrous citric acid, D&C
yellow no. 10, edetate disodium,
FD&C blue no. 1, flavors,
glycerin (soy), propylene glycol,
sodium benzoate, sorbitol,
sucralose, water, xanthan gum

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Made in England

Drug Facts**Active ingredients (in each 20 mL) Purposes**
MUCINEX FAST-MAX COLD & FLU ARCTIC BURST

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Drug Facts (continued)**Active ingredients (in each 20 mL) Purposes**
MUCINEX NIGHTSHIFT SEVERE COLD & FLU ARCTIC BURST

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine

Drug Facts (continued)**Uses MUCINEX FAST-MAX COLD & FLU ARCTIC BURST**

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

MUCINEX NIGHTSHIFT SEVERE COLD & FLU ARCTIC BURST

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sneezing
 - sinus congestion and pressure
 - runny nose
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have liver disease, heart disease, diabetes, high blood pressure, thyroid disease.

- glaucoma (Nightshift Severe Cold & Flu Arctic Burst only)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (Nightshift Severe Cold & Flu Arctic Burst only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
 - taking sedatives or tranquilizers
- (Nightshift Severe Cold & Flu Arctic Burst only)

When using this product do not use more than directed

- excitability may occur, especially in children (Nightshift Severe Cold & Flu Arctic Burst only)
- marked drowsiness may occur (Nightshift Severe Cold & Flu Arctic Burst only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nightshift Severe Cold & Flu Arctic Burst only)
- avoid alcoholic drinks (Nightshift Severe Cold & Flu Arctic Burst only)
- use caution when driving a motor vehicle or operating machinery (Nightshift Severe Cold & Flu Arctic Burst only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions**MUCINEX FAST-MAX COLD & FLU ARCTIC BURST**

- do not take more than directed (see Overdose warning)
 - do not take more than 6 doses in any 24-hour period
 - measure only with dosing cup provided
 - do not use dosing cup with other products
 - dose as follows or as directed by a doctor
 - adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
 - children under 12 years of age: do not use
- MUCINEX NIGHTSHIFT SEVERE COLD & FLU ARCTIC BURST**
- do not take more than directed (see Overdose warning)
 - do not take more than 4 doses in any 24-hour period
 - measure only with dosing cup provided
 - do not use dosing cup with other products
 - dose as follows or as directed by a doctor
 - adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
 - children under 12 years of age: do not use

Do not take Mucinex Fast-Max Cold & Flu Arctic Burst and Mucinex Nightshift Severe Cold & Flu Arctic Burst at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid.

Patents: www.b.com/patents

3181773 04/21/21 ©2021 RB Health

Phase visit our website www.mucinex.com

6382499546

5

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to the phone number
Maximum Strength per 4-hour dose
FAST-MAX COLD & FLU ARCTIC BURST helps to
relieve these symptoms day or night.

HEALTH • HYGIENE • HOME

MAXIMUM STRENGTH

Mucinex
FAST-MAX NIGHTSHIFT

DAY COLD & FLU **NIGHT** SEVERE COLD & FLU

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

FAST-MAX NIGHTSHIFT

MAXIMUM STRENGTH

DAY COLD & FLU **NIGHT** SEVERE COLD & FLU

Maximum Strength per 4-hour dose
Keep carton for full information.

PARENTS:
Learn about their medicine at
www.StopMedicineAbuse.org

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

MAXIMUM STRENGTH NDC 72854-241-20

Mucinex
FAST-MAX NIGHTSHIFT

DAY
COLD & FLU

Acetaminophen – Pain Reliever/
Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

NIGHT
SEVERE COLD & FLU

Acetaminophen – Pain Reliever/
Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant
Triprolidine HCl – Antihistamine



ALL IN ONE*

HEADACHE, SORE THROAT, FEVER, CHEST CONGESTION, COUGH, BODY PAIN, NASAL CONGESTION, SINUS CONGESTION, SINUS PRESSURE



ALL IN ONE*

HEADACHE, SORE THROAT, FEVER, ITCHY THROAT, COUGH, BODY PAIN, NASAL CONGESTION, RUNNY NOSE, SNEEZING

FAST RELEASE

12 CAPLETS **TOTAL 20 CAPLETS** **8 CAPLETS**

ACTUAL SIZE FOR AGES 12+ ACTUAL SIZE FOR AGES 12+

Questions?
1-866-MUCINEX (1-866-682-4639)
You may report side effects to
this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health

HEALTH • HYGIENE • HOME
www.mucinex.com
Patents: www.rb.com/patents

3176344 012721



6382499531 1

LOT:

EXP:

MADE IN:

3176344

Do not take the Mucinex Fast-Max Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. TAKE ONLY AS DIRECTED.

Drug Facts	Purposes
Active ingredients (in each caplet)	

010621 - - - - PEEL CORNER JO READ COMPLETE DRUG FACTS AND INFORMATION

Drug Facts (continued)

Do not use

- Ask a doctor before use if you have**

- Ask a doctor or pharmacist before use if you are**

- ### When using this product

- do not use more than directed
- excitability may occur, especially in children (*Nightshift Night Severe Cold & Flu only*)
- marked drowsiness may occur (*Nightshift Night Severe Cold & Flu only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Nightshift Night Severe Cold & Flu only*)
- avoid alcoholic drinks (*Nightshift Night Severe Cold & Flu only*)
- use caution when driving a motor vehicle or operating machinery (*Nightshift Night Severe Cold & Flu only*)

Hinge

Hinge

Drug Facts (continued)**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions**Mucinex Fast-Max Day Cold & Flu**

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Mucinex Nightshift Night Severe Cold & Flu

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information ■ store at 20-25°C (68-77°F)**Inactive ingredients Mucinex Fast-Max Day Cold & Flu**

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients Mucinex Nightshift Night Severe Cold & Flu

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.